


precautions<sup>2</sup> and call MCDPH to discuss

need for EVD testing

## Ebola Virus Disease (EVD) Evaluation Algorithm MCDPH: (602) 747-7111



Assessing Contact Exposure

## HIGH-RISK EXPOSURE

Does patient meet **ANY** of following within **21** days before symptom onset?

- -Percutaneous or mucous membrane exposure or direct skin contact with body fluids of a person with a confirmed or suspected case of EVD without appropriate personal protective equipment (PPE),
- -Laboratory worker processing body fluids of confirmed EVD patients without appropriate PPE or standard biosafety precautions
- -Participation in funeral rites which include direct exposure to human remains in the geographic area where outbreak is occurring without appropriate PPE

## LOW-RISK EXPOSURE

Does patient meet **ANY** of following within **21** days before symptom onset?
-Providing patient care (without known high-risk exposure) or contact with EVD patients in health care facilities in outbreak-affected countries<sup>1</sup>?
-Household member or close contact<sup>3</sup> of an EVD patient?

## **NO KNOWN EXPOSURE**

Does patient meet the following within **21** days before symptom onset? -Visited outbreak-affected countries<sup>1</sup>

<sup>2</sup>See CDC website for clinical, infection control and laboratory guidance and list of currently affected countries: www.cdc.gov/ebola

<sup>3</sup> Close contact is defined as a) being within approximately 3 feet (1 meter) or within the room or care area for a prolonged period of time (e.g., healthcare personnel, household members) while not wearing recommended personal protective equipment (i.e., droplet and contact precautions—see Infection Prevention and Control Recommendations); or b) having direct brief contact (e.g., shaking hands) with an EVD case while not wearing recommended personal protective equipment. At this time, brief interactions, such as walking by a person or moving through a hospital, do not constitute close contact. Symptoms include: Fever, weakness, muscle/joint pain, severe headache, stomach pain, vomiting, diarrhea, impaired kidney and liver function, unexplained bleeding or bruising, internal or external bleeding